For Research Use Only

EAAT4 Polyclonal antibody

Catalog Number: 12876-1-AP 2 Publications



Basic Information

Catalog Number: 12876-1-AP Concentration: 500 ug/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG3554

GenBank Accession Number:

BC028721 GeneID (NCBI): 6511 UNIPROT ID: P48664

solute carrier family 1 (high affinity aspartate/glutamate transporter),

Calculated MW: 61.6 kDa Observed MW: 60-70 kDa

Full Name:

member 6

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:2000

IF-Fro: 1:50-1:500

Positive Controls:

WB: A375 cells,

IF-Fro: mouse cerebellum tissue,

Applications

Tested Applications: WB, IF-Fro, ELISA Cited Applications: WB

Species Specificity: human, mouse, rat Cited Species: human, mouse

Background Information

Excitatory amino acid transporter 4 (EAAT4) is a neuronal glutamate transporter responsible for the reuptake of synaptically released glutamate. The EAAT4 protein is highly enriched in Purkinje cells of the cerebellum, although it is not restricted to these cells (18770868). While the predicted molecular weight of EAAT4 is 62 kDa, considerable heterogeneity in transporter size has been reported in the brain with molecular weights ranging from 50 to 120 kDa as a result of differential glycosylation and/or phosphorylation (24056007). This antibody recognizes 60-70 kDa of EAAT4 in mouse brain and several other lysate.

Notable Publications

Author	Pubmed ID	Journal	Application
Michael Rabenstein	29463856	Sci Rep	WB
Zhigang Huang	39733689	Neoplasia	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse cerebellum tissue using EAAT4 antibody (12876-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).

Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse cerebellum tissue using EAAT4 antibody (12876-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).

A375 cells were subjected to SDS PAGE followed by western blot with 12876-1-AP (EAAT4 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.