For Research Use Only

CD3 Polyclonal antibody Catalog Number:17617-1-AP Featured Product

Featured Product





Basic Information	Catalog Number: GenBank Accession Number: 17617-1-AP BC049847		Purification Method: Antigen affinity purification		
	Concentration:GeneID (NCBI):850 ug/ml916Source:ENSEMBL Gene ID:		:	Recommended Dilutions: WB: 1:1000-1:8000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Rabbit	ENSG0000019	8851	IHC: 1:500-1:2000	
	Isotype: IgG	UNIPROT ID: P07766 Full Name: CD3e molecule, epsilon (CD3-TCR complex)		IF-P: 1:50-1:500 IF-Fro: 1:200-1:800	
	Immunogen Catalog Number: AG11797				
		Calculated MV 207 aa, 23 kDa			
		Observed MW: 23-25 kDa			
Applications	Tested Applications: WB, IHC, IF-P, IF-Fro, IP, ELISA	Positive Cor			
	Cited Applications: mou: WB, IHC, IF, Cell treatment IP: Jr Species Specificity: IHC: human, mouse, rat, pig human Cited Species: human			WB : Jurkat cells, pig spleen tissue, MOLT-4 cells, mouse thymus tissue, rat thymus tissue IP : Jurkat cells,	
			-		
				IHC : human tonsillitis tissue, mouse spleen tissue, human lymphoma tissue, human lung cancer tissue, human breast cancer tissue, mouse colon tissue, human pancreas cancer tissue, rat thymus tissue IF-P : human tonsillitis tissue, mouse spleen tissue	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF-P : human		
				e colon tissue,	
Background Information	CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T- lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bour to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. This anti-CD3 is a polyclonal antibody directed against the epsilon cha of human CD3 molecule.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Xiao Wu	36168732	Exp Dermatol	IHC	
	Lan Zhang	34586738	Clin Transl Med	IHC,FC	
	Rilin Deng	32963356	Cell Mol Immunol	IHC	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol, pH7	7.3		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells (CD3+), MOLT-4 cells (CD3+), U-937 cells (CD3-), Raji cells (CD3-), mouse thymus tissue, and rat thymus tissue were subjected to SDS PAGE followed by western blot with 17617-1-AP (CD3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



WB result of CD3 antibody (17617-1-AP; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD3 transfected Jurkat cells.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 antibody (17617-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CD20 antibody (60271-1-Ig, Clone: 4A7G3, red).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 antibody (17617-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CD20 antibody (60271-1-Ig, Clone: 4A7G3, red).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17617-1-AP (CD3 antibody) at dilution of 1:1000 (under 10x (ens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17617-1-AP (CD3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CD3 (IP:17617-1-AP, 4ug; Detection:17617-1-AP 1:1000) with Jurkat cells lysate 3600ug.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse colon tissue using CD3 antibody (17617-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).