

For Research Use Only

PD-L1/CD274 Polyclonal antibody

Catalog Number:17952-1-AP

Featured Product

192 Publications



Basic Information

Catalog Number:

17952-1-AP

Concentration:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12432

GenBank Accession Number:

BC074984

GeneID (NCBI):

29126

UNIPROT ID:

Q9NZQ7

Full Name:

CD274 molecule

Calculated MW:

290 aa, 33 kDa

Observed MW:

45-56 kDa, 65-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:300-1:1200

IF/ICC 1:10-1:100

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, CHIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human placenta tissue, A375 cells, IFN gamma treated A549 cells, HeLa cells, HepG2 cells, human heart tissue, K-562 cells, mouse heart tissue, mouse skeletal muscle tissue, rat kidney tissue

IP: mouse heart tissue,

IHC: human tonsillitis tissue, mouse heart tissue, human stomach cancer tissue

IF/ICC: HEK-293 cells,

Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

Notable Publications

Author	Pubmed ID	Journal	Application
Ching-Yao Yang	29036791	Cancer Biomark	IHC
Gagan Chhabra	34597611	J Invest Dermatol	IHC
Haoyu Guo	36201949	Biomaterials	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human placenta tissue were subjected to SDS PAGE followed by western blot with 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Untreated and IFN gamma treated A549 cells were subjected to SDS PAGE followed by western blot with 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HEK-293 cells using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-PD-L1/CD274 (IP:17952-1-AP, 4ug; Detection:66248-1-Ig 1:1000) with mouse heart tissue lysate 4000ug.