

For Research Use Only

AXIN2 Polyclonal antibody

Catalog Number: 20540-1-AP

Featured Product

48 Publications



Basic Information

Catalog Number:

20540-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG14574

GenBank Accession Number:

BC006295

GeneID (NCBI):

8313

UNIPROT ID:

Q9Y2T1

Full Name:

axin 2

Calculated MW:

843 aa, 94 kDa

Observed MW:

94-100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : A549 cells, MDA-MB-453s cells, BxPC-3 cells, HeLa cells

Background Information

Axis inhibition protein2 (AXIN2), also known as Cductin or Axil, is a multidomain scaffold protein that negatively regulate Wnt signaling. AXIN2 can directly interact with beta-catenin and GSK3B. AXIN2 has a number of phosphorylation sites, and also can undergo poly(ADP-ribose)lation by tankyrase TNKS and TNKS2. Poly(ADP-ribose)lated AXIN2 then get ubiquitinated by RNF 146 and leads to its degradation and subsequent activation of Wnt signaling. AXIN2 is localized in the cytoplasm and its mutation is involved in colorectal cancer. Catalgo#20540-1-AP is a rabbit polyclonal antibody raised against a 34 amino acid N-terminal of human AXIN2. Observed MW of AXIN2 is from 93-100 kDa (PMID: 11809809; 24607787).

Notable Publications

Author	Pubmed ID	Journal	Application
Di Cui	34586717	Adv Healthc Mater	WB
Cefan Zhou	32972302	Autophagy	WB
Pingping Zhu	30224759	Nat Cell Biol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20540-1-AP (AXIN2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.