## For Research Use Only

## Perilipin-2 Monoclonal antibody

Catalog Number: 60340-1-lg



**Basic Information** 

Catalog Number: 60340-1-lg

 Size:
 GeneID (NCBI):

 1000 ug/ml
 123

 Source:
 UNIPROT ID:

Mouse Q99541
Isotype: Full Name:

IgG1 adipose differentiation-related

Immunogen Catalog Number: protein

AG7643 Calculated MW:

48 kDa
Observed MW:
47-52 kDa

BC005127

GenBank Accession Number:

**Applications** 

Tested Applications: WB, IHC, ELISA Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Protein A purification CloneNo.:

1G5D4

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:200-1:8000

Positive Controls:

WB: human adipose tissue,

IHC: human renal cell carcinoma tissue, mouse brown adipose tissue, rat brown adipose tissue

## **Background Information**

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage

Storage:

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-lg (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brown adipose slide using 60340-1 Ig (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).