

IMMUNOSTAINING CULTURED CELLS

www.ptglab.com

All steps in this protocol are performed at room temperature unless otherwise indicated. For optimum staining, incubations should be carried out on a slow-moving rotary shaker unless the cell line being used is delicate (e.g. neuronal cells).

Recipes for solutions highlighted **bold** are provided following the protocol.

-
- 1. Fixation and permeabilization:**
 - Aspirate medium, wash cells seeded on clean glass cover slips briefly with **1X PBS**.
 - Fix the cells with 4% paraformaldehyde made fresh in **1X PBS** for 10 minutes. Rinse cover slips with **1X PBS** 3 times for 3 minutes each.
 - Permeabilize with 0.2% Triton X-100 made in **1X PBS** for 5 minutes. Rinse cover slips 3 times with 1X PBS for 3 minutes each.

 - 2. Blocking:**
 - Prepare a blocking solution of 5% normal serum in **1X PBS**. Select serum from the same species in which the secondary antibody was raised e.g. if the secondary antibody is goat anti-mouse, then goat serum should be selected for the blocking solution. Incubate the cells with the blocking solution for 1 hour. (Alternatively, use 1% BSA in **1X PBS** for blocking if the corresponding serum is not available.)

 - 3. Antibody incubations:**
 - Aspirate the blocking solution and apply primary antibody diluted in **antibody dilution buffer**. Set aside one cover slipper experimental condition for a negative control and incubate in antibody dilution buffer minus the primary antibody. Leave these incubations for 1 hour, or, alternatively, incubate overnight at 4°C. **Please Note: If incubating overnight, take measures to ensure the cover slips do not dry out.**
 - Wash cover slips with **1X PBS** 3 times for 3 minutes each.
 - Apply an appropriate fluorophore-conjugated secondary antibody diluted in **antibody dilution buffer** to the coverslips and incubate for 1 hour in a moist, dark environment. **Please Note: It is imperative that cover slips be kept in dark conditions as much as possible after the addition of fluorescent reagents.**
 - Wash cover slips with **1X PBS** 3 times for 3 minutes each.

 - 4. Mounting and visualization:**
 - Mount cover slips on microscope slides with Hydromount (National Diagnostics) containing DAPI (if desired) for nuclear staining.
 - Examine slides under a fluorescence microscope.

IMMUNOSTAINING CULTURED CELLS

www.ptglab.com

Solutions

1X PBS	In 1000 ml (final volume)
10 mM Na ₂ HPO ₄	1.42 g
1.8 mM NaH ₂ PO ₄	0.22 g
140 mM NaCl	8.18 g
2.68mM KCl	0.20 g
Adjust to pH 7.4	
Add ddH ₂ O to 1000 ml	

Antibody dilution buffer	In 20 ml (final volume)
1% BSA	0.2 g
Add 1X PBS to 20ml	

Related Proteintech Products

Product Name	Catalog Number	Size	Applications
FITC-conjugated AffiniPure goat anti-mouse Ig (G+L)	SA00003-1	100 µl	FC, IF
FITC-conjugated AffiniPure goat anti-rabbit Ig (G+L)	SA00003-2	100 µl	FC, IF
TRITC-conjugated AffiniPure goat anti-mouse Ig (G+L)	SA00007-1	100 µl	FC, IF
TRITC-conjugated AffiniPure goat anti-rabbit Ig (G+L)	SA00007-2	100 µl	FC, IF