For Research Use Only

c-Myc Polyclonal antibody

Catalog Number:10057-1-AP 11 Publications

Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 10057-1-AP			Purification Method: Antigen affinity purification
	Size: 270 µg/ml	GenelD (NCBI): 4609		
	Source: Rabbit	UNIPROT ID: P01106		
	Isotype: Full Name: IgG v-myc myelocytomatosis viral oncogene homolog (avian)			
	Calculated MW: 48 aa, 8 kDa		W:	
	Observed MW: 62-65 kDa, 50 kDa			
Applications	Tested Applications: ELISA			
	Cited Applications: WB			
	Species Specificity: human, mouse, rat			
	Cited Species: human, rat, mouse			
	MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC. The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]			
Background Information	that regulates transcription binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan	le progression, apoptosis of specific target genes. manner, yet also specifi nent and translocation of phomas, including Burki t protein of human MYC.T	and cellular transfo It seems to activate cally recognizes the f this gene have been tt lymphoma. This a 'he 50kDa band reco	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of
	that regulates transcription binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan	le progression, apoptosis of specific target genes. manner, yet also specifi nent and translocation of phomas, including Burki t protein of human MYC.T	and cellular transfo It seems to activate cally recognizes the f this gene have been tt lymphoma. This a 'he 50kDa band reco	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of
	that regulates transcription binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan MYC, while the other bands	le progression, apoptosis of specific target genes. manner, yet also specifi nent and translocation of phomas, including Burki protein of human MYC.T , between 60-70kDa, are	and cellular transfo It seems to activate ically recognizes the f this gene have been tt lymphoma. This a he 50kDa band reco the phosphorylated	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of form of MYC [PMID: 12189186]
	that regulates transcriptior binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan MYC, while the other bands	le progression, apoptosis of specific target genes. manner, yet also specifi nent and translocation of phomas, including Burki t protein of human MYC.T , between 60-70kDa, are Pubmed ID	and cellular transfo It seems to activate ically recognizes the f this gene have been tt lymphoma. This a he 50kDa band reco the phosphorylated Journal	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of form of MYC [PMID: 12189186] Application
	that regulates transcription binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan MYC, while the other bands Author Chen Xin-Mei XM	le progression, apoptosis of specific target genes. manner, yet also specifi nent and translocation of phomas, including Burki t protein of human MYC.T , between 60-70kDa, are Pubmed ID 24265759	and cellular transfo It seems to activate ically recognizes the f this gene have been tt lymphoma. This a he 50kDa band reco the phosphorylated Journal PLoS One	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of form of MYC [PMID: 12189186] Application WB
Background Information	that regulates transcription binds DNA in a non-specific overexpression, rearranger tumors, leukemias and lym raised against recombinan MYC, while the other bands Author Chen Xin-Mei XM X Yuan	le progression, apoptosis of specific target genes. : manner, yet also specifi nent and translocation of iphomas, including Burki t protein of human MYC.T , between 60-70kDa, are Pubmed ID 24265759 27819668 32186775	and cellular transfo It seems to activate ically recognizes the f this gene have been tt lymphoma. This a he 50kDa band reco the phosphorylated Journal PLoS One Oncogenesis Mol Med Rep	rmation. It functions as a transcription factor the transcription of growth-related genes. It core sequence 5'-CAC[GA]TG-3'. Mutations, n associated with a variety of hematopoietic ntibody is a rabbit polyclonal antibody gnized by antibody is the native form of form of MYC [PMID: 12189186] Application WB WB

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer. Selected Validation Data