

PAFAH2 Polyclonal antibody

Catalog Number: 10085-1-AP

3 Publications

Basic Information

Catalog Number:

10085-1-AP

Size:

160 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0124

GenBank Accession Number:

BC001158

GeneID (NCBI):

5051

UNIPROT ID:

Q99487

Full Name:

platelet-activating factor
acetylhydrolase 2, 40kDa

Calculated MW:

40 kDa

Observed MW:

44-46 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse

Cited Species:

human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : mouse liver tissue,

IP : mouse liver tissue,

IHC : human lymphoma tissue,

Background Information

The family of platelet-activating factor-acetylhydrolase (PAF-AH) consists of enzymes which catalyze the hydrolysis of acetyl ester at the sn-2 position of PAF (Platelet-activating factor). The intracellular type II PAF-AH is a single polypeptide about a 40 kD distributed in both the cytosol and membranes of many kinds of cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Latchoumycandane Calivarathan C	22433871	J Biol Chem	WB
Thévenin Anastasia F AF	21882811	Biochemistry	WB, IF
Chen Jiawei J	17693621	J Lipid Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

**This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.**


Selected Validation Data



mouse liver tissue were subjected to SDS PAGE followed by western blot with 10085-1-AP (PAFAH2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 10085-1-AP (PAFAH2 antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-PAFAH2 (IP:10085-1-AP, 4ug; Detection:10085-1-AP 1:300) with mouse liver tissue lysate 6400ug.