For Research Use Only

SOX10 Polyclonal antibody

Catalog Number: 10422-1-AP

16 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10422-1-AP
 BC002824

 Size:
 GeneID (NCBI):

 800 µg/ml
 6663

Source: UNIPROT ID: Rabbit P56693
Isotype: Full Name:

gG SRY (sex determining region Y)-box

Immunogen Catalog Number: 1

AG0673 Calculated MW:

50 kDa Observed MW: 50-60 kDa

Applications

Tested Applications: FC, IHC, IP, WB, ELISA Cited Applications: CoIP, IF, IHC, WB Species Specificity:

human, mouse, rat Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: mouse brain tissue, mouse small intestine tissue,

Purification Method:

WB 1:1000-1:8000

protein lysate

IHC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

SH-SY5Y cells

IP: mouse colon tissue,
IHC: human brain tissue,

Background Information

Sox10 is a transcription factor that has a crucial role in complex signalling pathways regulating central nervous system and neural crest development. The calcualted molecular weight of SOX10 is 50 kDa, but modified SOX10 is about 55-60 kDa. (PMID: 23799842)

Notable Publications

Author	Pubmed ID	Journal	Application
Fränze Progatzky	34671159	Nature	IHC
Gelei Xiao	36524115	Front Immunol	IF
Erge Zhang	32572506	J Mol Med (Berl)	IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 10422-1-AP (SOX10 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-SOX10 (IP:10422-1-AP, 4ug; Detection:10422-1-AP 1:2000) with mouse colon tissue lysate 4000 ug.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 10422-1-AP (SOX10 antibody) at dilution of 1:100 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 C6 cells were intracellularly stained with 0.4 ug Anti-Human SOX10 (10422-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgC(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).