For Research Use Only

IBA1 Polyclonal antibody

Catalog Number: 10904-1-AP

Featured Product

345 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10904-1-AP BC009474

Size: GeneID (NCBI):

700 μg/ml 199

 Source:
 ENSEMBL Gene ID:

 Rabbit
 ENSG00000204472

 Isotype:
 UNIPROT ID:

 IgG
 P55008

Immunogen Catalog Number: Full Name:

AG1363 allograft inflammatory factor 1

Calculated MW: 17 kDa Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:400-1:1600 IF 1:50-1:500

Applications

Tested Applications:

FC, IF/ICC, IF-P, IHC, IP, WB, ELISA

Cited Applications: WB, IP, IF, IHC Species Specificity: human, mouse, rat Cited Species:

human, rat, sheep, mouse, zebrafish, hamster, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: THP-1 cells, IP: THP-1 cells,

IHC : rat brain tissue, human tonsillitis tissue, human liver tissue, mouse brain tissue, human brain tissue

IF: mouse brain tissue, rat brain tissue

Background Information

IBA1 is a 143 amino acid cytoplasmic, inflammation response scaffold protein. It is constitutively expressed in monocytes and macrophages and is known to be involved in macrophage activation. It is a marker of activated macrophage. Expression of IBA1 is up-regulated in activated microglia following facial nerve axotomy, ischemia, and several brain diseases, thereby implicating it in the activated phenotypes of microglia.

Notable Publications

Author	Pubmed ID	Journal	Application
Yi-Na Zhang	36168082	Transl Stroke Res	IF
Anna Piotrowska	31616413	Front Immunol	
R Zajaczkowska	30267794	Pharmacol Biochem Behav	

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of IBA1 antibody (10904-1-AP; 1:1000; room temperature for 1.5 hours) with wild-type and IBA1 knockout THP-1 cells.




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using IBA1 antibody (10904-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




1X10^6 THP-1 cells were intracellularly stained with 0.4 ug Anti-Human IBA1 (10904-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



IP result of anti-IBA1 (IP:10904-1-AP, 4ug; Detection:10904-1-AP 1:500) with THP-1 cells lysate 2400 ug.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).