For Research Use Only

Cofilin Polyclonal antibody

Catalog Number: 10960-1-AP

Featured Product

24 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10960-1-AP BC012318

Size: GeneID (NCBI): 1072

Source: UNIPROT ID: Rabbit P23528

Isotype: Full Name:
IgG cofilin 1 (non-muscle)
Immunogen Catalog Number: Calculated MW:

AG1399 18 kDa

Observed MW: 18-22 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:12000 IHC 1:20-1:200 IF 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB, IP, IF, IHC Species Specificity: human, mouse, rat Cited Species:

human, chicken, rat, mouse, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MCF-7 cells, NIH/3T3 cells, human skin tissue, human brain tissue, mouse brain tissue, rat brain tissue, NCI-H1299 cells

IHC: human cervical cancer tissue, human lung cancer tissue, human heart tissue

IF: NIH/3T3 cells,

Background Information

Cofilin is a ubiquitous actin-binding protein required for the reorganization of actin filaments. It is a member of ADF (actin-depolymerizing factor)/cofilin family that is a key regulator of actin dynamics and essential for cellular motility, cytokinesis, and endocytosis. Cofilin activity is tightly regulated by phosphorylation and dephosphorylation.Phosphorylation at Ser3 can inhibit its activity, also causing translocation from the nucleus to the cytoplasm.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhaoyue Meng	36175399	Nat Commun	WB
Kai Cui	27678468	Asian J Androl	WB
Jie Zan	27396619	Biochem Biophys Res Commun	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10960-1-AP (Cofilin antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human cervical cancer using 10960-1-AP (CFL1 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human cervical cancer using 10960-1-AP (CFL1 antibody) at dilution of 1:50 (under 40x lens)



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Cofilin antibody (10960-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).