For Research Use Only

RANGAP1 Polyclonal antibody

Catalog Number: 11102-1-AP

3 Publications



Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 11102-1-AP BC014044

Size: GeneID (NCBI): 5905

Source: UNIPROT ID: Rabbit P46060

Isotype: Full Name:

Ran GTPase activating protein 1

Immunogen Catalog Number: Calculated MW: AG1563 64 kDa

563 64 kDa

Observed MW:
64 kDa

Applications

Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, IP, WB Species Specificity: human

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SH-SY5Y cells, HEK-293 cells

IP: HEK-293 cells,

IHC: human gliomas tissue,

IF: HepG2 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Xianwei Chen	26844761	J Proteomics	WB
Hong-Juan You	37845585	Cell Oncol (Dordr)	WB,IF,CoIP
Yan Gong	36696903	Dev Cell	WB,IHC,IP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 11102-1-AP (RANGAP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11102-1-AP (RANGAP1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11102-1-AP (RANGAP1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 11102-1-AP (RANGAP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HepG2 cells were stained with 0.2ug RANGAP1 antibody (11102-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



IP result of anti-RANGAP1 (IP:11102-1-AP, 4ug; Detection:11102-1-AP 1:300) with HEK-293 cells lysate 3600ug.