## For Research Use Only

## FH Polyclonal antibody

Catalog Number: 11375-1-AP

12 Publications



**Basic Information** 

Catalog Number: 11375-1-AP

 Size:
 GeneID (NCBI):

 450 μ g/ml
 2271

Source: UNIPROT ID:
Rabbit P07954
Isotype: Full Name:
IgG fumarate hydratase

Immunogen Catalog Number:

AG1934 510 aa, 55 kDa

Observed MW: 45-50 kDa

Calculated MW:

BC003108

GenBank Accession Number:

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:250-1:1000 IF 1:50-1:500

**Applications** 

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

**Cited Applications:** 

IHC, WB

Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse, Caenorhabditis elegans

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells, human placenta tissue, HepG2 cells, LO2 cells, NIH/3T3 cells, mouse liver tissue, rat liver tissue, Hela cells, Jurkat cells, C6 cells

IP: mouse liver tissue,

IHC: human placenta tissue, human ovary tumor tissue

IF : HeLa cells,

**Background Information** 

Fumarate hydratase, or fumarase, is an enzymatic component of the tricarboxylic acid cycle, catalyzing the conversion of fumarate to malate. It belongs to the class-II fumarase/aspartase family. This protein has 2 isoforms produced by alternative initiation.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Teresa W-M Fan	36150727	J Immunol	
Yongyan Sun	30307357	Int J Radiat Biol	WB
Xiaoyu Ma	25301052	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 11375-1-AP (FH antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using FH antibody (11375-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-FH (IP:11375-1-AP, 4ug; Detection:11375-1-AP 1:8000) with mouse liver tissue lysate 3040 ug.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 11375-1-AP (FH antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 11375-1-AP (FH antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).