For Research Use Only

SIX2 Polyclonal antibody

Catalog Number: 11562-1-AP

Featured Product

245 Publications

BC024033

10736

Q9NPC8 Full Name:

GeneID (NCBI):

UNIPROT ID:

SIX homeobox 2 Calculated MW:

291 aa, 32 kDa

GenBank Accession Number:



Basic Information

Catalog Number: 11562-1-AP

Size: 750 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG2124

Observed MW: 32-37 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications:

IF, IP, ELISA

Cited Applications: ChIP, IP, WB

Species Specificity:

human **Cited Species:**

human, rat, mouse, rabbit, pig, Alligator, canine

Positive Controls:

IP: HEK-293 cells,

Background Information

The SIX proteins (sine oculis) are a family of homeodomain transcription factors that share a conserved DNA binding domain. Six members (Six1-Six6) of the Six gene family have been identified in mice and humans. SIX2, containing one homeobox DNA-binding domain, is highly expressed in fetal tissues but expression is limited in adult tissues. ${\it SIX2 may be involved in limb tendon and ligament development [PMID: 21420949]. It has been previously shown}$ that SIX2 is expressed in developing mesenchymal tissue including head and urogenital system at the time of overt midfacial and renal differentiation[PMID: 22282599]. This antibody is a rabbit polyclonal antibody raised against full length SIX2 of human origin.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------------|-----------|------------|-------------|
| Ravian L van Ineveld | 36180532 | Nat Protoc | |
| Bingjue Li | 32715474 | J Pathol | |
| Ken Hiratsuka | 36129975 | Sci Adv | |


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11562-1-AP (SIX2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 11562-1-AP (SIX2 antibody) at dilution of 1:5000 incubated at room temporarure for 1.5 hours.




IP result of anti-SIX2 (IP:11562-1-AP, 4ug; Detection:11562-1-AP 1:1000) with HEK-293 cells lysate 1320 ug.



"g", / Imministrationscend stating of SSL in Amount of SS

IF result from Fogelgren B, PMID:18570229, "Misexpression of Six2 is associated with heritable frontonasal dysplasia and renal hypoplasia in 3H1 Br mire"



IF result of SIX2 antibody (11562-1-AP, 1:200) with mouse embryonic kidney rudiment dissected at £13.5 and cultured for 2 days by Dr. Aleksandra Rak-Raszewska. SIX2 positive cells (red) incondense metanephric mesenchyme surrounding the ureteric bud tip.