For Research Use Only

BAFF/TNFSF13B Polyclonal antibody

Catalog Number: 11862-1-AP



Basic Information

Catalog Number:

11862-1-AP

BC020674

Concentration:

600 ug/ml

50urce:

Rabbit

Q9Y275

Isotype:

GenBank Accession Number:

BC020674

GeneID (NCBI):

10673

UNIPROT ID:

Q9Y275

Full Name:

IgG tumor necrosis factor (ligand)
Immunogen Catalog Number: superfamily, member 13b

AG2440 Calculated MW:

32 kDa Observed MW: 32 kDa, 45-52 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:2000 IHC: 1:200-1:800

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: U-937 cells, mouse testis tissue, mouse liver tissue, human liver tissue, rat liver tissue

IHC: human intrahepatic cholangiocarcinoma tissue,

Background Information

TNFSF13B, also known as BAFF, BLYS, TALL1, TNFSF20, ZTNF4 and CD257, is a cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. This cytokine is expressed in B lineage cells, and acts as a potent B cell activator. The BAFF system promotes B cell survival and differentiation and plays a prominent role in the pathogenesis of autoimmune diseases. It has been shown that BAFF inhibitors are in clinical trials for systemic lupus erythematosus with significant efficacy. This antibody recognizes all the forms of TNFSF13B such as soluble form (17 kDa), membrane form (32-35 kDa) and glycosylation form (45-52 kDa).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



U-937 cells were subjected to SDS PAGE followed by western blot with 11862-1-AP (TNFSF 13B antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 11862-1-AP (TNFSF13B antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 11862-1-AP (TNFSF13B/BAFF antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).