

For Research Use Only

# NDUFS1 Polyclonal antibody

Catalog Number: 12444-1-AP

Featured Product

70 Publications



## Basic Information

### Catalog Number:

12444-1-AP

### Size:

450 µg/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG3135

### GenBank Accession Number:

BC030833

### GeneID (NCBI):

4719

### UNIPROT ID:

P28331

### Full Name:

NADH dehydrogenase (ubiquinone)

Fe-S protein 1, 75kDa (NADH-coenzyme Q reductase)

### Calculated MW:

727 aa, 79 kDa

### Observed MW:

74-81 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:250-1:1000

IF 1:200-1:800

## Applications

### Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

### Cited Applications:

WB, IF, IHC, ColP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, rat, mouse, zebrafish, Drosophila

### Positive Controls:

WB: mouse kidney tissue, mouse liver tissue, HEK-293 cells, A549 cells, HeLa cells, HEK-293 tissue, rat kidney tissue

IP: mouse lung tissue,

IHC: human liver cancer tissue,

IF: HepG2 cells,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

The multisubunit NADH:ubiquinone oxidoreductase (75 kDa subunit, mitochondrial)(NDUFS1) is the first enzyme complex in the electron transport chain of mitochondria. It is also named as Complex I-75kD. By use of chaotropic agents, complex I can be fragmented into 3 different fractions: a flavoprotein fraction, an iron-sulfur protein (IP) fraction, and a hydrophobic protein (HP) fraction. NDUFS1 is the largest subunit of complex I and it is a component of the iron-sulfur (IP) fragment of the enzyme. NDUFS1 has some isoforms with the molecular mass of 68-81 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Shao-Ming Shen	26415504	EMBO Rep	WB
Yang Ni	31557978	Cells	WB
Rufeng Zhang	31530015	FASEB J	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12444-1-AP (NDUFS1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




WB result of NDUFS1 antibody (12444-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NDUFS1 transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 12444-1-AP (NDUFS1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 12444-1-AP (NDUFS1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12444-1-AP (NDUFS1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12444-1-AP (NDUFS1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NDUFS1 antibody (12444-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



IP result of anti-NDUFS1 (IP:12444-1-AP, 4ug; Detection:12444-1-AP 1:500) with mouse lung tissue lysate 2800ug.