For Research Use Only

FAS/CD95 Polyclonal antibody

Catalog Number: 13098-1-AP

Featured Product

114 Publications

BC012479

GeneID (NCBI):

UNIPROT ID:

P25445 Full Name:

GenBank Accession Number:

Fas (TNF receptor superfamily,



Basic Information

Catalog Number: 13098-1-AP

Size: 1000 µg/ml

Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG3718

35-38 kDa Observed MW:

38-45 kDa

member 6)

Calculated MW:

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications: FC, IP, WB, ELISA

Cited Applications: WB, IP, IF, IHC, CoIP, CHIP

Species Specificity:

human Cited Species:

human, sheep, monkey, hamster

Positive Controls:

WB: HT-1080 cells, HT-29 cells, Jurkat cells, HeLa cells

IP: HeLa cells,

Background Information

Fas (CD95/APO-1) is a transmembrane glycoprotein belonging to the tumor necrosis factor (TNF) receptor superfamily. It can mediate apoptosis by ligation with an agonistic anti-Fas antibody or Fas ligand. Stimulation of Fas results in the aggregation of its intracellular death domains, leading to the formation of the death-inducing signaling complex (DISC). FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. The molecular mass of native Fas is 38 kDa, the high molecular weight form (40-55 kDa) of Fas is due to glycosylation.

Notable Publications

Author	Pubmed ID	Journal	Application
Shifeng Pan	29152131	Oncotarget	WB
Lan Zhang	34586738	Clin Transl Med	WB
Xinghuo Wu	31554783	Exp Mol Med	WB,IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of FAS/CD95 antibody (13098-1-AP, 1:1000) with si-control and si-FAS transfected Jurkat cells.



Various lysates were subjected to SDS PAGE followed by western blot with 13098-1-AP (FAS/CD95 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-FAS/CD95 (IP:13098-1-AP, 3ug; Detection:13098-1-AP 1:500) with HeLa cells lysate 3000ug.



1x10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human FAS/CD95 (13098-1-AP) (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).