### For Research Use Only

# Aggrecan Polyclonal antibody

Catalog Number: 13880-1-AP

225 Publications



**Basic Information** 

Catalog Number: 13880-1-AP

GenBank Accession Number:

BC036445

Size: 500 μg/ml GeneID (NCBI):

**UNIPROT ID:** 

Rabbit P16112 Full Name: Isotype: aggrecan Calculated MW: Immunogen Catalog Number:

250 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

IHC 1:200-1:800

**Applications** 

**Tested Applications:** 

FC, IHC, ELISA

Cited Applications:

IF, IHC

Source:

Species Specificity:

human, mouse, rat

**Cited Species:** 

human, rat, mouse, rabbit, zebrafish, pig, canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

IHC: human lung tissue, human lung cancer tissue, human cartilage tissue, mouse lung tissue

# **Background Information**

Aggrecan (ACAN), also named as CSPCP, AGC1 and MSK16, belongs to the aggrecan/versican proteoglycan family (also including versican, brevican and neurocan). It is a major component of extracellular matrix of cartilaginous tissues. A major function of aggrecan is to resist compression in cartilage. It binds avidly to hyaluronic acid via an Nterminal globular region.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yangfei Yi	36178080	Orthop Surg	IHC
Sa Cha	34638972	Int J Mol Sci	IHC
Lindsay N Collins	31571216	J Comp Neurol	


#### Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# Selected Validation Data



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 13880-1-AP (Aggrecan antibody at dilution of 1:400 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 13880-1-AP (Aggrecan antibody at dilution of 1:400 (under 40x lens).



1X10^6 ATDC-5 cells were intracellularly stained with 0.8 ug Anti-Human Aggrecan (13880-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).