## For Research Use Only

# CRISP3 Polyclonal antibody

Catalog Number: 14847-1-AP

Featured Product

**4 Publications** 



**Basic Information** 

Catalog Number: 14847-1-AP

Size:

Source:

Rabbit

650 μg/ml

GenBank Accession Number:
NM\_006061
GeneID (NCBI):
10321
UNIPROT ID:
P54108

 Isotype:
 Full Name:

 IgG
 cysteine-rich secretory protein 3

Calculated MW: 28 kDa Observed MW: 29 kDa **Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications:

IHC, IP, WB, ELISA
Cited Applications:

IHC, WB

Species Specificity: human, mouse Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human saliva, human placenta tissue, Jurkat cells, mouse colon tissue, mouse ovary tissue, mouse thymus tissue

IP: human placenta tissue,

IHC: human prostate cancer tissue, human pancreas tissue

# **Background Information**

CRISP3, also known as SGP28, is a glycoprotein that belongs to a family of cysteine-rich secretory proteins (CRISPs). CRISP3 was originally discovered in human neutrophils, but transcripts are widely distributed in exocrine glands (salivary glands, pancreas, and prostate) and also found at lower levels in epididymis, ovary, thymus, and colon. CRISP3 is a potential biomarker for prostate cancer. This antibody can recognize both CRISP2 and CRISP3.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yu-Ming Wang	36263172	Int J Biol Sci	WB,IHC
XLiu	36267015	J Dent Res	WB
Paola A Rojas	28376177	J Natl Cancer Inst	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




human saliva samples were subjected to SDS PAGE followed by western blot with 14847-1-AP (CRISP3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 14847-1-AP (CRISP3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 14847-1-AP (CRISP3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CRISP3 (IP:14847-1-AP, 4ug; Detection:14847-1-AP 1:300) with human placenta tissue lysate 1520ug.