For Research Use Only

Connexin-43 Polyclonal antibody

Catalog Number: 15386-1-AP 34 Publications



Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:50-1:200

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 15386-1-AP BC026329 GeneID (NCBI): Size: 330 μg/ml 2697 **UNIPROT ID:** Source: Rabbit P17302 Isotype: Full Name:

gap junction protein, alpha 1, 43kDa

Calculated MW: Immunogen Catalog Number: AG6698 43 kDa

> Observed MW: 43-47 kDa

Applications

Tested Applications: FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, IP, WB

Species Specificity: human, mouse, rat **Cited Species:**

human, rat, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls: WB: HeLa cells,

IP: mouse heart tissue,

IHC: mouse heart tissue, human heart tissue

IF: HeLa cells, mouse heart tissue

Background Information

Connexin-43 (Cx43, also known as gap junction alpha-1), a member of connexin family, plays essential roles in gap junction communication that facilitates direct communication among adjacent cells (PMID: 12270943). Usually, six connexin proteins oligomerize into a hemi-channel or connexon during intercellular channel formation (PMID: 12270943). Mutations of Cx43 may cause a series of diseases such as oculodentodigital dysplasia (ODDD), syndactyly 3 (SDTY3), hypoplastic left heart syndrome 1 (HLHS1) and so on (PMID: 18161618, 1472936, 1147490).

Notable Publications

Author	Pubmed ID	Journal	Application
Xueling He	28969971	Prog Biophys Mol Biol	WB
E O Adegoke	30267989	Ecotoxicol Environ Saf	IF
Hsiang-Chun Lee	28935953	Sci Rep	WB,IP

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse heart using 15386-1-AP (GJA1 antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded mouse heart using 15386-1-AP (GJA1 antibody) at dilution of 1:100 (under 40x lens).




1X10^6 Jurkat cells were stained with 0.5ug Connexin-43 antibody (15386-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



HeLa cells were subjected to SDS PAGE followed by western blot with 15386-1-AP (GJA1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-Connexin-43 (IP:15386-1-AP, 4ug; Detection:15386-1-AP 1:500) with mouse heart tissue lysate 3200ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 15386-1-AP (Connexin-43 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).