

FCGR2A / CD32a Polyclonal antibody

Catalog Number: 15625-1-AP

9 Publications

Basic Information

Catalog Number:

15625-1-AP

Size:

550 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8072

GenBank Accession Number:

BC020823

GeneID (NCBI):

2212

UNIPROT ID:

P12318

Full Name:

Fc fragment of IgG, low affinity IIa, receptor (CD32)

Calculated MW:

35 kDa

Observed MW:

35-40 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:1000-1:4000

Applications

Tested Applications:

FC, IHC, WB, ELISA

Cited Applications:

WB, IF, FC, IHC

Species Specificity:

human

Cited Species:

human, rat, mouse

Positive Controls:

WB : K-562 cells, U-937 cells

IHC : human liver tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IgG Fc receptors (FcγRs) play important roles in immune responses. Low affinity IgG Fc receptor Fc gamma RIIa (FCGR2A, CD32a) is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenlong Zhao	36116152	Int Immunopharmacol	WB, IF
Luodan Yang	29056360	Exp Neurol	WB
Jianbin Yin	35739102	Cell Death Dis	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15625-1-AP (FCGR2A / CD32a antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 15625-1-AP (FCGR2A / CD32a antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 15625-1-AP (FCGR2A / CD32a antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ K-562 cells were intracellularly stained with 0.4 ug Anti-Human FCGR2A / CD32a (15625-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).