## For Research Use Only

## SLC3A1 Polyclonal antibody

Catalog Number:16343-1-AP

Featured Product







Basic Information	Catalog Number: GenBank Accession Number: 16343-1-AP BC022386		cession Number:	Purification Method: Antigen affinity purification Recommended Dilutions:	
	Size:	GenelD (NCBI):			
	350 µg/ml	6519		WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500	
	Source:	UNIPROT ID	t.		
	Rabbit	Q07837			
	lsotype: IgG	Full Name: solute carrier family 3 (cystine, dibasic and neutral amino acid			
	Immunogen Catalog Number:				
	AG9525	transporters, activator of cystine, dibasic and neutral amino acid transport), member 1 Calculated MW: 685 aa, 79 kDa Observed MW:			
					79 kDa
	Applications	Tested Applications:		Positive Cor	ntrols:
IHC, IP, WB, ELISA			WB : HEK-293 cells, mouse kidney tissue, rat kidne		
Cited Applications: IF, IHC, WB			tissue		
Species Specificity:			idney tissue,		
numan, mouse, rat		IHC : human tissue	uman kidney tissue, human small intestine		
Cited Species:					
goat, human, mouse					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	SLC3A1(Neutral and basic amino acid transport protein rBAT) is also named as NBAT,D2h,RBAT. It is involved in the high-affinity, sodium-independent transport of cystine and neutral and dibasic amino acids (system B(0,+)-like activity).SLC3A1 can be degraded, at least in part, via the endoplasmic reticulum-associated degradation (ERAD) pathway(PMID:18332091). SLC3A1 has different isoforms vary from 79 kDa to 36kDa and the 85 kDa glycosylated protein can be detected in SDS-PAGE. (PMID:11546643)				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Gangqi Wang	36332571	Cell Stem Cell	IF	
			Cai Dan		
	Lauren E Woodard	28317878	Sci Rep	IF	
	Lauren E Woodard Lauren E Woodard	28317878 31221135	BMC Nephrol	IF WB,IF	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 16343-1-AP (SLC3A1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16343-1-AP (SLC3A1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). IP result of anti-SLC3A1 (IP:16343-1-AP, 4ug; Detection:16343-1-AP 1:1000) with mouse kidney tissue lysate 4000ug.