

# ATP1A2 Polyclonal antibody

Catalog Number: 16836-1-AP

16 Publications

## Basic Information

## Catalog Number:

16836-1-AP

## Size:

700 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG10515

## GenBank Accession Number:

BC052271

## GeneID (NCBI):

477

## UNIPROT ID:

P50993

## Full Name:

ATPase, Na<sup>+</sup>/K<sup>+</sup> transporting, alpha 2  
(+) polypeptide

## Calculated MW:

1020 aa, 112 kDa

## Observed MW:

97-100 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:10-1:100

## Applications

## Tested Applications:

FC, IF/ICC, IHC, WB, ELISA

## Cited Applications:

IF, IHC, WB

## Species Specificity:

human, mouse, rat

## Cited Species:

canine, *Haliotis discus hannai*, human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

**WB** : 37°C incubated mouse heart tissue, 37°C incubated mouse skeletal muscle tissue

**IHC** : mouse heart tissue, human kidney tissue, human testis tissue, human skin tissue, human heart tissue

**IF** : HeLa cells,

## Background Information

ATP1A2 (Na<sup>+</sup>/K<sup>+</sup>-ATPase α-2 subunit) is the catalytic component of the active enzyme Na<sup>+</sup>/K<sup>+</sup>-ATPase, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. The Na<sup>+</sup>/K<sup>+</sup>-ATPase is composed of a larger catalytic α-subunit (~110 kDa) and a small β-subunit (~55 kDa). The α subunit has four isoforms identified to date: α 1, α 2, α 3 and α 4. The α 1 isoform is expressed ubiquitously but the α 2 isoform is present largely in the skeletal muscle, heart and vascular smooth muscle. The α 3 isoform is found almost exclusively in neurons and ovaries. The α 4 isoform is expressed in sperm. This antibody was raised against the internal region of the human ATP1A2 and can recognize all the isoforms of α subunit. The 65kDa band detected occasionally may be the degradation product of ATP1A2.

## Notable Publications

Author	Pubmed ID	Journal	Application
Ji Zhu	28970012	Eur J Pharmacol	WB
Yanglei Jia	30245637	Front Physiol	WB
Mariarosaria Cammarota	34481380	Biomed Pharmacother	WB, IF

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16836-1-AP (ATP1A2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




37 °C incubated or boiled mouse skeletal muscle lysates were subjected to SDS PAGE followed by western blot with 16836-1-AP (ATP1A2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 16836-1-AP (ATP1A2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HeLa cells, using ATP1A2 antibody 16836-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



1X10<sup>6</sup> HeLa cells were stained with 0.5ug ATP1A2 antibody (16836-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). FITC-Goat anti-Rabbit IgG with dilution 1:100.