

IFT52 Polyclonal antibody

Catalog Number: 17534-1-AP

Featured Product

22 Publications

Basic Information

Catalog Number:

17534-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11586

GenBank Accession Number:

BC039831

GeneID (NCBI):

51098

UNIPROT ID:

Q9Y366

Full Name:intraflagellar transport 52 homolog
(Chlamydomonas)**Calculated MW:**

437 aa, 50 kDa

Observed MW:

45-50 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IF 1:10-1:100

Applications

Tested Applications:

IF/ICC, IP, WB, ELISA

Cited Applications:

WB, IP, IF, ColP

Species Specificity:

human, mouse, rat, Canine

Cited Species:

human, Chicken, rat, mouse

Positive Controls:**WB** : HEK-293 cells, human testis tissue, mouse testis
tissue, MDCK cells, LNCaP cells**IP** : mouse testis tissue, HeLa cells**IF** : HeLa cells,

Background Information

IFT52, also named as C20orf9 and NGD5, is a component of IFT complex B which is composed of IFT88, IFT57, TRAF3IP1, IFT52, IFT27, HSPB11 and IFT20. IFT52 binds directly to the IFT81/74/27/25 complex. And IFT52 could mediate the interaction between the two subcomplexes. (PMID:21642430) The MW of IFT52 is about 45-54kd.

Notable Publications

Author	Pubmed ID	Journal	Application
Ivan Duran	27666822	Sci Rep	WB
Shichao Duan	33241915	EMBO J	WB
Chunyu Liu	31735294	Am J Hum Genet	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17534-1-AP (IFT52 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HeLa cells, using IFT52 antibody 17534-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-IFT52 (IP:17534-1-AP, 4ug; Detection:17534-1-AP 1:1000) with mouse testis tissue lysate 4000ug.