For Research Use Only

Cytokeratin 10 Polyclonal antibody

Catalog Number: 18343-1-AP

13 Publications



Basic Information

Catalog Number: 18343-1-AP Size:

 $600 \mu g/ml$ Source: Rabbit Isotype: Immunogen Catalog Number:

AG13136

GenBank Accession Number:

BC034697 GeneID (NCBI): 3858

P13645 Full Name: keratin 10 Calculated MW:

UNIPROT ID:

584 aa, 59 kDa

Observed MW: 50-59 kDa

Positive Controls:

WB: mouse skin tissue, A431 cells, rat skin tissue

Purification Method:

WB 1:5000-1:20000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

IHC: human cervical cancer tissue, mouse skin tissue, rat skin tissue, human lung cancer tissue, human breast cancer tissue, human skin cancer tissue

Applications

Tested Applications: FC, IHC, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse, rat **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. As a type I keratin, keratin 10 is a suprabasal marker of differentiation in stratified squamous epithelia.

Notable Publications

Author	Pubmed ID	Journal	Application
Lisa Lemoine	34579573	mBio	IF
Yingzheng Zhao	36240893	Int J Biol Macromol	IF
Shidi Wu	31790685	Life Sci	IHC

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:200000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human Cytokeratin 10 (18343-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).