

CAPG-Specific Polyclonal antibody oroteintech Antibodies | ELISA kits | Proteins Catalog Number: 19535-1-AP www.ptglab.com Featured Product **1** Publications Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 19535-1-AP NM_001747 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 240 µg/ml 822 WB 1:500-1:2000 IHC 1:20-1:200 UNIPROT ID: Source: Rabbit P40121 Full Name: Isotype: IgG capping protein (actin filament), gelsolin-like Calculated MW: 39 kDa **Observed MW:** 43 kDa **Applications Tested Applications:** Positive Controls: IHC, WB, ELISA WB : mouse lung tissue, mouse kidney tissue, U-937 **Cited Applications:** cells WB IHC : human spleen tissue, human stomach cancer Species Specificity: tissue human, mouse, rat **Cited Species:** human Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Background Information** CapG (gelsolin-like actin-capping protein) is a member of the gelsolin superfamily proteins, which regulates actin filament length by capping or severing filaments. Studies have shown a role for CapG in regulating cell motility. The expression of CapG is upregulated under hypoxia and in turn increases the motility of cells. Recently it has been reported that the overexpression of CapG is associated with the progression of lung adenocarcinoma as well as lyphatic invasion status of cholangiocarcinoma, which may be used for prognosis of these tumors. (21908955,22155129) Notable Publications Author Pubmed ID Journal Application Shao Fangchun F WB 21908955 Tohoku J Exp Med Storage: Storage Store at -20°C. Stable for one year after shipment. Storage Buffer PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage


For Research Use Only

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







mouse lung tissue were subjected to SDS PAGE followed by western blot with 19535-1-AP (CAPG-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human spleen using 19535-1-AP (CAPG-Specific antibody) at dilution of 1:50 (under 10x lens). Immunohistochemical analysis of paraffinembedded human spleen using 19535-1-AP (CAPG-Specific antibody) at dilution of 1:50 (under 40x lens).