### For Research Use Only

# RIG-1/DDX58 Polyclonal antibody

Catalog Number: 20566-1-AP

Featured Product

41 Publications



**Basic Information** 

Catalog Number: 20566-1-AP

1000 µg/ml Source: Rabbit

Isotype: IgG

Size:

GenBank Accession Number:

NM\_014314 GeneID (NCBI): 23586 UNIPROT ID: 095786 Full Name:

DEAD (Asp-Glu-Ala-Asp) box

polypeptide 58
Calculated MW:
107 kDa
Observed MW:
101/106 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF 1:10-1:100

**Applications** 

Tested Applications: IF/ICC, IP, WB, ELISA Cited Applications: CoIP, IHC, IP, RIP, WB Species Specificity: human, mouse, rat Cited Species:

human, mouse, monkey, pig

**Positive Controls:** 

WB: Jurkat cells, A431 cells, MCF-7 cells, mouse brain

tissue, HepG2 cells, MOLT-4 cells

IP: HepG2 cells,

IF: HepG2 cells, H9C2 cells

# **Background Information**

DDX58, also named as RIG-1, belongs to the helicase family. It is involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAVS/IPS1, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). Detects dsRNA produced from non-self dsDNA by RNA polymerase III, such as Epstein-Barr virus-encoded RNAs (EBERs). It is essential for the production of interferons in response to RNA viruses including paramyxoviruses, influenza viruses, Japanese encephalitis virus and HCV. The antibody is specific to DDX58.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Liang Zhang	30258002	J Virol	WB
Wei Zhang	25228491	J Gen Virol	WB
Lei-Ke Zhang	27605671	J Virol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 20566-1-AP (RIG-1/DDX58 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using DDX58 antibody 20566-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



WB result of RIG-1/DDX58 antibody (20566-1-AP; 1:2000; room temperature for 1.5 hours) with wild-type and RIG-1/DDX58 knockout A431 cells.



IP result of anti-RIG-1/DDX58 (IP:20566-1-AP, 4ug; Detection:20566-1-AP 1:300) with HepG2 cells lysate 3600ug.