For Research Use Only

AIM2 Polyclonal antibody

Catalog Number: 20590-1-AP

Featured Product

21 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 20590-1-AP
 BC010940

 Size:
 GeneID (NCBI):

 750 µg/ml
 9447

 Source:
 UNIPROT ID:

 Rabbit
 014862

Rabbit O14862
Isotype: Full Name:
IgG absent in melanoma 2

Immunogen Catalog Number:Calculated MW:AG14520343 aa, 39 kDa

Observed MW: 39-42 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000 IHC 1:20-1:200

Applications

Tested Applications: IHC, WB, ELISA Cited Applications: IHC, WB

Species Specificity: human, mouse, rat Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HeLa cells, Jurkat cells, SH-SY5Y cells, mouse testis tissue, rat testis tissue, MCF-7 cells, A549 cells, COLO 320 cells, HepG2 cells, Raji cells

IHC: human testis tissue,

Background Information

Absent in melanoma 2 (AIM2) is a sensor of cytosolic DNA that is responsible for activation of the inflammasome and host immune responses to DNA viruses and intracellular bacteria [PMID: 22695634]. Upon binding to DNA, it is thought to undergo oligomerization and to associate with PYCARD initiating the recruitment of caspase-1 precusrsor and processing of interleukin-1 beta and interleukin-18. Otherwise, it functions as a tumor suppressor which may act by repressing NF-kappa-B transcriptional activity [PMID:16432157,20566831].

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------------------|-----------|--------------------------|-------------|
| Liangmin Fu | 36248876 | Front Immunol | WB,IHC |
| JiangFan Yu | 36248980 | Front Oncol | WB |
| Saleela M Ruwanpura | 36037355 | Proc Natl Acad Sci U S A | WB,IHC |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20590-1-AP (AIM2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 20590-1-AP (AIM2 antibody) at dilution of 1:100 (under 10x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 20590-1-AP (AIM2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.