For Research Use Only

GAD65 Polyclonal antibody

Catalog Number:20746-1-AP

14 Publications




Basic Information	Catalog Number: 20746-1-AP	GenBank Accession Number: NM_000818		Purification Method: Antigen affinity purification	
	Size:	GenelD (NC		Recommended Dilutions: WB 1:3000-1:20000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	750 µg/ml	2572			
	Source:	UNIPROT ID):		
	Rabbit	Q05329		protein lysate IF 1:10-1:100	
	Isotype: Full Name: IgG glutamate decarboxylase 2 (pancreatic islets and brain, 65kD				
	Calculated MW: 65 kDa				
	Observed MW: 65 kDa				
Applications	Tested Applications:		Positive Controls: WB : mouse brain tissue, rat brain tissue		
	IF/ICC, IP, WB,ELISA				
	Cited Applications: IF, IHC, WB	IP : mouse brain tissue,			
	Species Specificity: IF : HeLa cells, human, mouse, rat				
	Cited Species: human, rat, mouse				
Background Information	GAD2, also named as GAD65, belongs to the group II decarboxylase family. GAD2 catalyzes the production of GAB/ It is responsible for the synthesis of the essential neurotransmitter gamma-aminobutyric acid (GABA) from L- glutamic acid. GAD2 is expressed in nervous and endocrine systems and are thought to be involved in synaptic transmission and INS secretion. Autoantibodies against GAD2 may serve as markers for type I diabetes. Many individuals suffering from an adult onset disorder known as Stiff Person Syndrome (SPS) also express autoantibodies to GAD2. The antibody is specific to GAD2.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Yueqin Liu	33344711	Neurobiol Stress	WB	
	Hongru Yang	35618610	Adv Sci (Weinh)	WB	
	Ken Chen	31015358	Clin Sci (Lond)	IHC	
Storage	Storage: Store at -20°C. Stable for or Storage Buffer:	ne year after shipment.			


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20746-1-AP (GAD65 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HeLa cells using 20746-1-AP (GAD65 antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-GAD65 (IP:20746-1-AP, 4ug: Detection:20746-1-AP 1:1000) with mouse brain tissue lysate 3440ug.