

TTC19 Polyclonal antibody

Catalog Number: 20875-1-AP **2 Publications**

Basic Information

Catalog Number: 20875-1-AP	GenBank Accession Number: BC011698	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 54902	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q6DKK2	
Isotype: IgG	Full Name: tetra-tricopeptide repeat domain 19	
Immunogen Catalog Number: AG14974	Calculated MW: 501 aa, 56 kDa	
	Observed MW: 32-35 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : mouse heart tissue, HeLa cells, rat heart tissue IHC : human breast cancer tissue, mouse heart tissue
Cited Applications: IHC, WB	
Species Specificity: human, mouse, rat	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

TTC19 is a subunit of mitochondrial respiratory chain complex III and is proposed to preserve the structural and functional integrity of complex III. Loss-of-function mutations in TTC19 have been found in patients with complex III deficiency associated with heterogeneous neurological syndromes. The precursor TTC19 was detected as a 43 kDa cytoplasmic protein, while the mature TTC19 with the mitochondrial targeting sequence cleaved was observed as 35 kDa by SDS-PAGE. (PMID: 21278747)

Notable Publications

Author	Pubmed ID	Journal	Application
Huijuan Dai	37928256	Int J Biol Sci	WB,IHC
Aojia Zhuang	37158593	Elife	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20875-1-AP (TTC19 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 20875-1-AP (TTC19 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)).