

For Research Use Only

Cytokeratin 14 Polyclonal antibody

Catalog Number: 22221-1-AP

1 Publications



Basic Information

Catalog Number:

22221-1-AP

Size:

300 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG17559

GenBank Accession Number:

BC002690

GeneID (NCBI):

3861

UNIPROT ID:

P02533

Full Name:

keratin 14

Calculated MW:

472 aa, 52 kDa

Observed MW:

52 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

FC, IF-P, IHC, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, hamster

Positive Controls:

WB : mouse skin tissue, rat skin tissue

IP : mouse skin tissue,

IHC : human skin tissue, human lung cancer tissue, human breast cancer tissue, human cervical cancer tissue

IF : mouse skin tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral counterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. Keratin 14 is a type I cytokeratin. It is usually found as a heterotetramer with keratin 5. Keratins K14 and K5 have long been considered to be biochemical markers of the stratified squamous epithelia, including epidermis. This antibody is specifically against KRT14.

Notable Publications

Author	Pubmed ID	Journal	Application
S Vijayalingam	24874842	Cancer Gene Ther	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 22221-1-AP (Cytokeratin 14 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human skin tissue slide using 22221-1-AP (Cytokeratin 14 antibody at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human skin tissue slide using 22221-1-AP (Cytokeratin 14 antibody at dilution of 1:200 (under 40x lens).




IP result of anti-Cytokeratin 14 (IP:22221-1-AP, 4ug; Detection:22221-1-AP 1:1000) with mouse skin tissue lysate 3200ug.



1X10⁶ A431 cells were intracellularly stained with 0.2 ug Anti-Human Cytokeratin 14 (22221-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Control Antibody. Cells were fixed with 90% MeOH.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skin tissue using Cytokeratin 14 antibody (22221-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skin tissue using Cytokeratin 14 antibody (22221-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).