## For Research Use Only

## TBP Polyclonal antibody

Catalog Number:22246-1-AP

2 Publications



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 22246-1-AP
 BC110341

 Size:
 GeneID (NCBI):

 300 μ g/ml
 6908

 Source:
 UNIPROT ID:

 Rabbit
 P20226

Isotype: Full Name:
IgG TATA box binding protein

Immunogen Catalog Number:Calculated MW:AG17740338 aa, 38 kDaObserved MW:

mouse/rat 33-36 kDa and human 37-43kDa

**Applications** 

Tested Applications: IF/ICC, WB,ELISA Cited Applications: WR

Species Specificity: human, mouse Cited Species:

human, mouse

**Positive Controls:** 

WB: Jurkat cells, COLO 320 cells, HeLa cells, mouse colon tissue, SKOV-3 cells

**Purification Method:** 

WB 1:500-1:2000 IF 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

IF: MCF-7 cells,

Background Information

TBP, TATA box-binding protein, is a DNA-binding subunit of TFIID, which is a multisubunit complex essential for the expression of most protein-encoding genes. The well-conserved C-terminal domains of 180 aa is both necessary and sufficient for interaction with DNA and for assembly of the basal transcription apparatus. The N terminus of TBP modulates the DNA-binding activity of the C terminus of the protein. Once binds to the TATA box mediated by TBP, TFIID initiate the transcription step of the pre-initiation complex (PIC), playing a role in the activation of eukaryotic genes transcribed by RNA polymerase II. This is a rabbit polyclonal antibody raised against part of N-terminal chain of human TBP, and specially recognize a ~46 kDa band.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Qingye Li	27748821	Mol Med Rep	WB
Hiroyuki Minemura	33713004	Breast Cancer	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 22246-1-AP (TBP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of MCF-7 cells, using TBP antibody 22246-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).