For Research Use Only

CSNK2B Polyclonal antibody

Catalog Number:22418-1-AP

2 Publications



Basic Information

Catalog Number:

22418-1-AP

Size:

450 µg/ml

Source:

Rabbit

BC112017

GeneID (NCBI):

1460

UNIPROT ID:

Rabotype:

Full Name:

casein kinase 2, beta polypeptide

Immunogen Catalog Number:Calculated MW:AG18134215 aa, 25 kDaObserved MW:

25-30 kDa

Applications

Tested Applications: FC, IHC, WB, ELISA Cited Applications: WB

Species Specificity: human, mouse Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, HepG2 cells, NIH/3T3 cells, K-562

Purification Method:

WB 1:1000-1:6000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cells, MCF-7 cells

IHC: human testis tissue, human thyroid tissue, mouse testis tissue, mouse brain tissue

Background Information

CSNK2B is a ubiquitous protein kinase which regulates metabolic pathways, signal transduction, transcription, translation, and replication. The enzyme is composed of three subunits, alpha, alpha prime and beta, which form a tetrameric holoenzyme. The alpha and alpha prime subunits are catalytic, while the beta subunit serves regulatory functions. The enzyme localizes to the endoplasmic reticulum and the Golgi apparatus. It participates in Wnt signaling, and plays a complex role in regulating the basal catalytic activity of the alpha subunit.

Notable Publications

Author	Pubmed ID	Journal	Application
Hiroki Okanishi	36357940	Cancer Metab	WB
Yan Zhang	31987502	Biochem Biophys Res Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 22418-1-AP (Casein Kinase 2 beta antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 22418-1-AP (Casein Kinase 2 beta antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 22418-1-AP (Casein Kinase 2 beta antibody at dilution of 1:200 (under 40x lens).



1X10^6 HeLa cells were stained with 0.2ug Casein Kinase 2 beta antibody (22418-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.