For Research Use Only

CEP89, CCDC123 Polyclonal antibody



Purification Method:

WB 1:200-1:1000 IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Catalog Number: 24002-1-AP

3 Publications

Basic Information

Applications

Catalog Number: 24002-1-AP Size: 220 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG21206

85-100 kDa, 40 kDa

Tested Applications: IF/ICC, WB, ELISA

Cited Applications:

WB. IF

Species Specificity: human, mouse **Cited Species:** human, mouse

GenBank Accession Number:

BC136328 GeneID (NCBI): 84902 **UNIPROT ID:** Q96ST8

Calculated MW:

783 aa, 90 kDa Observed MW:

Full Name: coiled-coil domain containing 123

Positive Controls:

WB: SH-SY5Y cells, Neuro-2a cells

IF: U-251 cells,

Background Information

CCDC123(as known as CEP123), also named as CEP89, is a new player in the process of primary ciliogenesis and it also plays a role in mitochondrial metabolism where it may modulate complex IV activity. It has been shown that CEP123 is localized to the distal appendages of the mother centriolecep and the localization of CEP123 is cell cycledependent with its levels decreasing during mitosis. CEP123 depletion can cause defects in ciliary vesicle formationcep and prevent the formation of a ciliary vesicle at the distal end of the mother centriole. It is possible that CEP123 is involved in regulating the recruitment of membranes to the centrosome through its interaction with Cep290(PMID:23575228, 23789104, 23348840). 24002-1-AP antibody recognizes all of CEP123 isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Pierrick Le Borgne	36070319	PLoS Biol	IF
Min Liu	34480124	Cell Res	WB,IF
Tomoharu Kanie	28625565	Dev Cell	IF

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

Selected Validation Data



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 24002-1-AP (CEP89 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.

Neuro-2a cells were subjected to SDS PAGE followed by western blot with 24002-1-AP (CEP89 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using 24002-1-AP (CEP89 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).