

NDNL2 Polyclonal antibody

Catalog Number: 27488-1-AP

1 Publications

Basic Information

Catalog Number:

27488-1-AP

Size:

230 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26663

GenBank Accession Number:

BC053999

GeneID (NCBI):

56160

UNIPROT ID:

Q96MG7

Full Name:

necdin-like 2

Calculated MW:

34 kDa

Observed MW:

38 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, L02 cells, HeLa cells, human placenta tissue

IHC : human liver cancer tissue,

IF : HepG2 cells,

Background Information

NDNL2 (also known as NSMCE3 or MAGEG1) is a part of the SMC5-6 protein complex that is essential for DNA damage response and chromosome segregation (PMID: 18086888; 27427983). The gene of human NDNL2 maps to chromosome 15q13.1. Northern blot analysis detected expression of a 1.9-kb transcript in all human tissues tested, with highest expression in testis (PMID: 11782285). Missense mutations in NSMCE3 have been associated with an autosomal recessive chromosome breakage syndrome that leads to defective T and B cell function and acute respiratory distress syndrome in early childhood (PMID: 27427983).

Notable Publications

Author	Pubmed ID	Journal	Application
Stephanie Pei Tung Yiu	35263599	Cell Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2, L02 and HeLa cells were subjected to SDS PAGE followed by western blot with 27488-1-AP (NDNL2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27488-1-AP (NDNL2 antibody) at dilution of 1:100 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27488-1-AP (NDNL2 antibody) at dilution of 1:100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 27488-1-AP (NDNL2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).