For Research Use Only

ORAI1 Polyclonal antibody

Catalog Number:28411-1-AP

Featured Product

5 Publications





Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 28411-1-AP BC015369 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 300 µg/ml 84876 WB 1:500-1:1000 IHC 1:50-1:500 UNIPROT ID: Source: Rabbit Q96D31 Full Name: Isotype: lgG ORAI calcium release-activated calcium modulator 1 Immunogen Catalog Number: AG29234 Calculated MW: 301 aa. 33 kDa **Observed MW:** 45-55 kDa **Applications Tested Applications: Positive Controls:** IHC, WB, ELISA WB: MDA-MB-453s cells, MCF-7 cells **Cited Applications:** IHC : mouse testis tissue, WB, IF Species Specificity: Human, mouse **Cited Species:** bovine, human, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Background Information** Store-operated calcium channels (SOCs) play essential functions in various physiological processes. ORAI1 is a complex glycosylated protein, it exists in several molecular weight forms: a theoretical molecular weight of 32.7 kDa of the unmodified protein and two major glycosylated proteins of ~43 kDa and ~50 kDa (PMID: 22318387). Human patients with a severe combinaed immunodeficiency (SCID) syndrome possess point mutations in Orai1 gene. **Notable Publications** Author Pubmed ID Journal Application Qingjie Kang 34055617 Front Oncol WB J Anim Sci IF Ming Li 34100951 WB.IF **Cheng-Shun Zhang** Life Sci 34102196 Storage Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





MDA-MB-453s cells were subjected to SDS PAGE
followed by western blot with 28411-1-AP (ORAI1
antibody) at dilution of 1:500 incubated at room
temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 28411-1-AP (ORAI1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

MCF-7 cells were subjected to SDS PAGE followed by western blot with 28411-1-AP (ORAI 1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.