For Research Use Only

SARS-CoV-2 Nucleocapsid Phosphoprotein Polyclonal antibody



Catalog Number: 28769-1-AP

2 Publications

Basic Information

Catalog Number:

28769-1-AP

600 µg/ml

Source:

Rabbit Isotype:

laC.

Size:

Immunogen Catalog Number:

AG30676

GenBank Accession Number:

NC_045512 GeneID (NCBI):

43740575 Full Name:

COVID-19 N Protein

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

Virus

Cited Species:

human

Positive Controls:

WB: Recombinant protein,

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. it can be used as a marker in diagnostic assays due to its high immunogenicity.(PMID: 32416961, PMID: 32235387)

Notable Publications

Author	Pubmed ID	Journal	Application
Yi Zheng	35075101	Signal Transduct Target Ther	WB
Yongxuan Yao	36464077	Antiviral Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



SARS-CoV-2 Nucleocapsid Phosphoprotein with the domain 1-419aa expressed in HEK-293 cells were subjected to SDS PAGE followed by western blot with 28769-1-AP (2019-nCOV nucleocapsid phosphoprotein antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



SARS-CoV-2 Nucleocapsid Phosphoprotein
Antibody (28769-1-AP) tested by ELISA.SARS-CoV-2
Nucleocapsid Phosphoprotein was coated onto
microtiter plates at 0.15 µg/well and then
incubated with a dilution series of SARS-CoV-2
Nucleocapsid Phosphoprotein Antibody (28769-1AP). Bound antibodies were detected with HRP
conjugated anti-Rabbit IgG followed by incubation
with HRP Substrate and then measuring the
resulting absorbance at 450 nm.