For Research Use Only

GDI1 Polyclonal antibody

Catalog Number:55214-1-AP




| Basic Information | Catalog Number: 55214-1-AP | GenBank Accession Number: NM_001493 | Purification Method: Antigen affinity purification | |
|---|---|--|---|--------------|
| | Size: 800 µg/ml | GenelD (NCBI): 2664 | Recommended Dilutions: WB 1:500-1:2000 | |
| | Source: Rabbit | UNIPROT ID: P31150 | | |
| | lsotype: IgG | Full Name: GDP dissociation inhibitor 1 | | |
| | | Calculated MW: 51 kDa Observed MW: 51-55 kDa | | |
| | | | | Applications |
| WB,ELISA Species Specificity: human, mouse, rat | WB : SH-SY5Y cells, human brain tissue, mouse brain tissue | | | |
| Background Informatio | family. GDIs can bind and rel proteins have been identifier not Rab3A. GDI1 is expressed diffuse, cytoplasmic distribut Defects in GDI1 are the cause | GDP dissociation inhibitors (GDIs) are proteins that regulate the GDP-GTP exchange reaction of members of the rab family. GDIs can bind and release GDP-bound Rab proteins from membranes. Two GDI proteins towards different Ra proteins have been identified. GDI1 interacts with almost all of the Rab proteins, while GDI2 interacts with Rabll bu not Rab3A. GDI1 is expressed primarily in neural and sensory tissues and also in secretory cells, displaying a diffuse, cytoplasmic distribution in cells. It runs as a 55kda protein in SDS-PAGE. (PMID: 7929030, PMID: 19570034). Defects in GDI1 are the cause of mental retardation X-linked type 41 (MRX41). Defects in GDI1 are the cause of mental retardation X-linked type 48 (MRX48). The antibody is specific to GDI1. And it has no cross reaction to GDI2. | | |
| Storage | Storage: Store at -20°C. Storage Buffer: PBS with 0.02% sodium azide Aliquoting is unnecessary for | 0, 1 | | |

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 55214-1-AP (GDI1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.