

FBXW7 Polyclonal antibody

Catalog Number: 55290-1-AP

Featured Product

10 Publications

Basic Information

Catalog Number:

55290-1-AP

Size:

700 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_018315

GeneID (NCBI):

55294

UNIPROT ID:

Q969H0

Full Name:

F-box and WD repeat domain containing 7

Calculated MW:

80 kDa

Observed MW:

100-110 kDa, 66-75 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, HEK-293 cells, MCF-7 cells, human brain tissue

IP : HepG2 cells,

IHC : human skin cancer tissue, human colon cancer tissue

IF : HeLa cells,

Background Information

FBXW7, also named as FBW7, FBX30, SEL10 and hAgo, is a substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. It probably recognizes and binds to phosphorylated target proteins. FBXW7 is involved in the degradation of cyclin-E, MYC, NOTCH1 released notch intracellular domain (NICD), and probably PSEN1. FBXW7 is a general tumor suppressor in human cancer (PMID: 17909001). FBXW7 has 3 isoforms (α / β / γ) with the calculated molecular mass of 80 kDa, 70 kDa and 66 kDa, and apparent molecular mass of 100-110 kDa and 66-75 kDa (PMID: 18559665). This antibody is specific to FBXW7 and it can recognize all the isoforms of FBXW7.

Notable Publications

Author	Pubmed ID	Journal	Application
Heling Dong	26360620	Endocrinology	WB
Haiwei Zhang	28507201	Biosci Rep	WB
Xia Li	31257502	Mol Med Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of FBXW7 antibody (55290-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-FBXW7 transfected HEK-293 cells.




HepG2 cell lysates were subjected to SDS PAGE followed by western blot with 55290-1-AP (FBXW7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human skin cancer tissue slide using 55290-1-AP (FBXW7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 55290-1-AP (FBXW7 antibody), at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-FBXW7 (IP:55290-1-AP, 4ug; Detection:55290-1-AP 1:500) with HepG2 cells lysate 960 ug.