## For Research Use Only

## SIRT1 Monoclonal antibody

Catalog Number:60303-1-lg Featured Product 76 Publications



## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 60303-1-lg BC012499 Protein A purification GenelD (NCBI): Concentration: CloneNo.: 2000 µg/ml 23411 7F6G6 UNIPROT ID: Recommended Dilutions: Source: Mouse Q96EB6 WB: 1:5000-1:50000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total Full Name: Isotype: protein lysate lgG2b sirtuin (silent mating type IHC: 1:250-1:1000 information regulation 2 homolog) 1 Immunogen Catalog Number: IF/ICC: 1:1000-1:4000 (S. cerevisiae) AG17677 Calculated MW: 747 aa, 82 kDa Observed MW: 110-130 kDa **Applications Tested Applications: Positive Controls:** WB, IHC, IF/ICC, IP, ELISA WB: A549 cells, HeLa cells, Hek-293 cells, LNCaP cells, **Cited Applications:** Jurkat cells, MCF-7 cells WB, IHC, IF, CoIP, ELISA IP: HeLa cells. **Species Specificity:** IHC : human breast cancer tissue, human testis tissue, human human colon tissue **Cited Species:** IF/ICC : BT-549 cells, human, pig, rabbit Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Background Information** SIRT1, also named as SIR2L1, contains a deacetylase sirtuin-type domain and belongs to the sirtuin family. The posttranslation modified SIRT1 is a 110-130 kDa protein, which contains one deacetylase sirtuin-type domain. The 75-80 kDa SIRT1 fragment was detected to lack the carboxy-terminus (PMID:21305533). SIRT1 exists a 57-61 kDa isoform. SIRT1 may be found in nucleolus, nuclear euchromatin, heterochromatin and inner membrane. It can shuttles between nucleus and cytoplasm. SIRT1 regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. SIRT1 in particular initiates several signaling events relevant to cardioprotection, including: activation of endothelial nitric oxide synthase, INS receptor signaling, and autophagy. In addition SIRT1 activation elicits resistance to oxidative stress via regulation of transcription factors and co-activators such as FOXO, Hif-2a, and NF-kB. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lysine 382. **Notable Publications** Author Pubmed ID Journal Application Jie Liu 36238648 Oxid Med Cell Longev WB Xin Peng 34549339 J Mol Neurosci WB J Orthop Surg Res ChuanCai Liang 36064420 WB Storage: Storage Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60303-1-lg (SIRT1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed BT-549 cells using SIRT1 antibody (60303-1-1g, Clone: 7F6G6) at dilution of 1:2000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).




WB result of SIRT1antibody (60303-1-lg, 1:500) with si-Control and si-SIRT1 transfected HEK-293 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 60303-1-lg (SIRT1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



IP result of anti-SIRT1 (IP:60303-1-Ig, 5ug; Detection:60303-1-Ig 1:1000) with HeLa cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60303-1-1g (SIRT1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).