

For Research Use Only

CHOP; GADD153 Monoclonal antibody



Catalog Number: 60304-1-Ig

11 Publications

Basic Information

Catalog Number:

60304-1-Ig

Size:

2000 µg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG7354

GenBank Accession Number:

BC003637

GeneID (NCBI):

1649

UNIPROT ID:

P35638

Full Name:

DNA-damage-inducible transcript 3

Calculated MW:

19 kDa

Observed MW:

30 kDa

Purification Method:

Protein A purification

CloneNo.:

4D5A9

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human

Cited Species:

human, rat, mouse

Positive Controls:

WB : Tunicamycin treated HeLa cells,

Background Information

CHOP, also known as GADD153 or DDIT3, is a highly conserved gene in both the structural and regulatory regions. Imposed by unfolded and misfolded proteins, CHOP is significantly induced by ER stress. CHOP is considered a proapoptotic marker of ER stress dependent cell death. CHOP acts as a dominant-negative inhibitor of the transcription factor C/EBP and LAP. It may play an important role in the malignant transformation of nevus to melanoma. The calculated molecular weight of CHOP is 19 kDa, but the protein migrates on an SDS-PAGE gel with an observed molecular mass of 29 kDa (PMID: 1547942).

Notable Publications

Author	Pubmed ID	Journal	Application
Yaoyao Bian	36121296	Pharm Biol	IF
Valeria Catena	27655709	Oncotarget	WB
Adrian Rivera-Reyes	30382078	Cell Death Dis	WB, IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Tunicamycin treated HeLa cells were subjected to SDS PAGE followed by western blot with 60304-1-Ig (CHOP; GADD153 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.