For Research Use Only

CD68 Monoclonal antibody

Catalog Number:66231-2-lg 71 Publications



Basic Information

Catalog Number: GenBank Accession Number: 66231-2-lg BC015557

Concentration: Genel D (NCBI): 2000 ug/ml 968

Source: UNIPROT ID:

Mouse P34810

Isotype: Full Name:
IgG1 CD68 molecule

Immunogen Catalog Number: Calculated MW:
AG22815 37 kDa

Purification Method: Protein G purification

CloneNo.:

3A9A7

Recommended Dilutions: IHC: 1:1000-1:6000 IF-P: 1:3000-1:12000

Applications

Tested Applications: IHC, IF-P, ELISA Cited Applications:

IHC, IF

Species Specificity:

human
Cited Species:
human, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue, human appendicitis tissue, human colon cancer tissue, human liver cancer tissue, human lung cancer tissue, human urothelial carcinoma tissue

IF-P: human tonsillitis tissue,

Background Information

CD68 is a type I transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It belongs to the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. CD68 is also a member of the scavenger receptor family. It may play a role in phagocytic activities of tissue macrophages.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Huang	34713723	J Am Heart Assoc	
Yourong Jiang	36389736	Front Immunol	IF
Xinyi Zhang	36499018	Int J Mol Sci	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-Ig, Clone: 3A9A7) at dilution of 1:6000 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-Ig, Clone: 3A9A7) at dilution of 1:6000 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-lg (CD68 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).