For Research Use Only

Cyclin A2 Monoclonal antibody

Catalog Number:66391-1-lg Featured Product

23 Publications




Basic Information	Catalog Number: 66391-1-Ig	GenBank Accession Numl BC104783	per: Purification Method: Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	1400 µg/ml	890	1C6C11
	Source:	UNIPROT ID:	Recommended Dilutions:
	Mouse	P20248	WB 1:2000-1:10000
	Isotype: IgG2a	Full Name: cyclin A2	IHC 1:1000-1:4000 IF 1:50-1:500
	Immunogen Catalog Number: AG12765	Calculated MW: 432 aa, 49 kDa	
		Observed MW: 52 kDa	
Applications	Tested Applications:	Pc	ositive Controls:
	IF/ICC, IHC, WB, ELISA Cited Applications:		B : HepG2 cells, HeLa cells, HEK-293 cells, Jurkat Ils, Raji cells
	IF, WB	IH	IC : human tonsillitis tissue,
	Species Specificity: human	IF	: HeLa cells, MCF-7 cells
	Cited Species: human, rat, mouse		
	Note-IHC: suggested an with TE buffer pH 9.0; (` antigen retrieval may b with citrate buffer pH 6.	*) Alternatively, e performed	
Background Information	in protein abundance through the distinct expression and degradat contrast to cyclin A1, which is pre	e cell cycle. Cyclins function as ion patterns which contribute t esent only in germ cells, this cy	embers are characterized by a dramatic periodicit regulators of CDK kinases. Different cyclins exhib o the temporal coordination of each mitotic event. clin is expressed in all tissues tested. CCNA2 bind ell cycle G1/S and G2/M transitions.
Notable Publications	Author	Pubmed ID Journal	Application
	Y Zhong	28869606 Oncogene	e WB
	Dianbao Zhang	33076708 Nutr Cano	er WB
	Jingwen Tan	36208777 Chem Bio	l Interact WB
Storage	Storage: Store at -20°C. Stable for one yea	r after chinment	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66391-1-1g (Cyclin A2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of Cyclin A2 antibody (66391-1-Ig; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Cyclin A2 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66391-1-Ig (Cyclin A2 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsilliits tissue slide using 66391-1-1g (Cyclin A2 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 66391-1-Ig(Cyclin A2 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).