For Research Use Only

AGTR1 Monoclonal antibody

Catalog Number:66415-1-lg 1 Publications




Basic Information	Catalog Number: 66415-1-lg	GenBank Accession Number: BC022447	Purification Method: Protein A purification	
	Size: 1300 µg/ml	GenelD (NCBI): 185	CloneNo.: 2C5G11	
	Source: Mouse	UNIPROT ID: P30556	Recommended Dilutions: WB 1:1000-1:4000	
	lsotype: IgG2a	Full Name: angiotensin II receptor, type :	1	
	Immunogen Catalog Number: AG14461	Calculated MW: 359 aa, 41 kDa		
		Observed MW: 50 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB,ELISA Cited Applications: WB	WB : pig heart tissue, pig skeletal muscle tissue		
	Species Specificity: human, rat, pig, mouse			
	Cited Species: mouse			
Background Information	Angiotensin II (Ang II), the main effector molecule of the renin-angiotensin system, exerts its actions mainly via interaction with type-1 angiotensin II receptor (AGTR1, also named as AT1R), thereby contributing to blood pressur regulation. AGTR1 mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. By regulating vascular tone, cardiovascular function, salt and water homeostasis, AGTR exerts an indispensable physiological role (PMID: 21600887). AGTR1 has been implicated in diverse aspects of human disease, from the regulation of blood pressure and cardiovascular homeostasis to cancer progression (PMID 26975580).			
Notable Publications	Author	Pubmed ID Journal	Application	
	Han Su	34603779 J Adv Res	WB	
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



pig heart tissue were subjected to SDS PAGE followed by western blot with 66415-1-1g (AGTR1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.