

# Caspase 3/p17/p19 Monoclonal antibody

Catalog Number: 66470-2-Ig

Featured Product

215 Publications

## Basic Information

Catalog Number:

66470-2-Ig

Size:

1500 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG25029

GenBank Accession Number:

BC016926

GeneID (NCBI):

836

UNIPROT ID:

P42574

Full Name:

caspase 3, apoptosis-related cysteine peptidase

Calculated MW:

277 aa, 32 kDa

Observed MW:

32-35 kDa, 19 kDa, 17 kDa

Purification Method:

Protein G purification

CloneNo.:

2G4B2

Recommended Dilutions:

WB 1:1000-1:3000

IHC 1:150-1:600

IF 1:50-1:500

## Applications

Tested Applications:

IF/ICC, IHC, WB, ELISA

Cited Applications:

ICC, IF, IHC, WB

Species Specificity:

Human, mouse

Cited Species:

human, chicken, rat, mouse, pig, plant, canine

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : Jurkat cells, HEK-293 cells, HepG2 cells, NIH/3T3 cells

IHC : human breast cancer tissue, mouse liver tissue, mouse kidney tissue

IF : human breast cancer tissue, HepG2 cells

## Background Information

Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning of apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19 and p32 of Caspase 3.

## Notable Publications

Author	Pubmed ID	Journal	Application
Jingjing Zheng	32978798	Ann N Y Acad Sci	WB
Yang Liu	36149580	Cell Stress Chaperones	WB
Yaling Zhang	36233452	J Clin Med	IF

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




WB result of Caspase 3 antibody (66470-2-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Caspase 3 transfected HEK-293 cells.




Untreated and Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 66470-2-Ig (CASP3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66470-2-Ig (CASP3 antibody) at dilution of 1:300 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66470-2-Ig (CASP3 antibody) at dilution of 1:300 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using 66470-2-Ig (CASP3 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Caspase 3/p17/p19 antibody (66470-2-Ig, Clone: 2G4B2) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).