For Research Use Only

## OLIG2 Monoclonal antibody, PBS Only




Catalog Number:66513-1-PBS

| Basic Information                         | Catalog Number:<br>66513-1-PBS  | GenBank Accession Number:<br>BC047511   | Purification Method:<br>Protein A purification |
|---|---|---|--|
|   | Size:<br>1 mg/ml<br>Source:<br>Mouse<br>Isotype:<br>IgG2a<br>Immunogen Catalog Number:<br>AG18838   | GenelD (NCBI):<br>10215   | CloneNo.:<br>1E8C5                             |
|   |   | UNIPROT ID:<br>Q13516   |  |
|   |   | Full Name:<br>oligodendrocyte lineage transcription<br>factor 2<br>Calculated MW:<br>32 kDa |  |
|   |   |   |  |
|   |   | Applications  | Tested Applications:<br>WB,ELISA               |
| Species Specificity:<br>Human, mouse, rat |   |   |  |
| Background Information                    | OLIG2, also named as BHLHB1, BHLHE19, PRKCBP2 and RACK17, is required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1, OLIG2 establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development. OLIG2 is widely expressed in subsets of glia cells and progenitors, and it is strongly induced at different sites by both acute and chronic injury, albeit with different mechanisms. OLIG2 acts as a repressor of neurogenesis in cells reacting to brain injury. It may represent an effective approach towards evoking neuronal repair from parenchymal precursors.(PMID:19390819) |   |  |
| Storage                                   | Storage:<br>Store at -20°C. Stable for one year<br>Storage Buffer:<br>PBS only<br>Aliquoting is unnecessary for -20°C   |   |  |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 66513-1-Ig (OLIG2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66513-1-PBS in a different storage buffer formulation.