For Research Use Only

FMR1 Monoclonal antibody

Catalog Number:66548-1-lg Featured Product






Basic Information	Catalog Number: 66548-1-lg	GenBank A BC038998	ccession Number:	Purification Method: Protein A purification	
	Size: 2100 µg/ml	Genel D (NO 2332	CBI):	CloneNo.: 2B5B5	
	Source: Mouse	UNIPROT II 006787):	Recommended Dilutions: WB 1:2000-1:10000	
	Isotype: IgG2a Immunogen Catalog Number: AG4697	Full Name: fragile X mental retardation 1 Calculated MW: 632 aa, 71 kDa		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:250-1:1000	
	Applications	Tested Applications:		Positive Co	Positive Controls:
WB, IHC, FC (Intra), IP, ELISA Cited Applications:				WB : HeLa cells, HEK-293 cells, K-562 cells, Jurkat cells, HSCT6 cells IP : HeLa cells,	
WB, IF			IP : HeLa ce		
human mouse rat			human gliomas tissue, mouse brain tissue, an liver tissue, human colon cancer tissue, humar		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Informatior	polymbosomes, implicating inte	egulation of trans IRNA. It also has a	lation. FMR1 is a compo a role in the transport of	protein complex that associates with onent of the CYFIP1-EIF4E-FMR1 complex f mRNA from the nucleus to the cytoplasm. 59-72 kDa.	
Notable Publications	Author	Pubmed ID	Journal	Application	
	Belinda J Petri	36341798	Environ Res	WB	
	Raeann Goering	32510328	Elife	WB	
	Cheng Wu	37152432	Front Mol Neurosc		
Storage	Storage: Store at -20°C. Stable for one yea	ar after chinment			


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66548-1-lg (FMR1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-FMR1 (IP:66548-1-Ig, 5ug; Detection:66548-1-Ig 1:3000) with HeLa cells lysate 3200 ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 66548-1-1g (FMR1 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human FMR1 (66548-1-1g, Clone:2B5B5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-1g, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).