

# EIF4E Monoclonal antibody

Catalog Number: 66655-1-Ig

Featured Product

2 Publications

## Basic Information

Catalog Number:

66655-1-Ig

Size:

1800 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG27191

GenBank Accession Number:

BC012611

GeneID (NCBI):

1977

UNIPROT ID:

P06730

Full Name:

eukaryotic translation initiation factor 4E

Calculated MW:

29 kDa

Observed MW:

26-29 kDa

Purification Method:

Protein A purification

CloneNo.:

3C6B9

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:2500-1:10000

IF 1:50-1:500

## Applications

Tested Applications:

FC, IF/ICC, IHC, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

Human, mouse, rat

Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

**WB** : Jurkat cells, U2OS cells, LNCaP cells, HeLa cells, HEK-293 cells, HSC-T6 cells, NIH/3T3 cells, RAW 264.7 cells, MCF-7 cells

**IHC** : human breast cancer tissue,

**IF** : HepG2 cells,

## Background Information

Eukaryotic translation initiation factor 4E, also known as eIF4E, is a protein that in humans is encoded by the EIF4E gene. eIF4E is the mRNA cap-binding protein, known as a general initiation factor allowing for mRNA-ribosome interaction and cap-dependent translation in eukaryotic cells. eIF4E is a polypeptide that exists as both a free form and as part of the eIF4F pre-initiation complex. Regulation of eIF4E may be achieved via three distinct mechanisms: transcription, phosphorylation, and inhibitory proteins.

## Notable Publications

Author	Pubmed ID	Journal	Application
Shasha Zhao	35768165	J Immunother Cancer	WB
Xiao Ke	35802246	Neurosci Bull	WB,IHC

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66655-1-Ig (EIF4E antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66655-1-Ig (EIF4E antibody) at dilution of 1:5000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 66655-1-Ig (EIF4E antibody) at dilution of 1:100 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



WB result of EIF4E antibody (66655-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EIF4E transfected HEK-293 cells.



1X10<sup>6</sup> HepG2 cells were intracellularly stained with 0.2 ug Anti-Human EIF4E (66655-1-Ig, Clone:3C6B9) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Various lysates were subjected to SDS PAGE followed by western blot with 66655-1-Ig (EIF4E antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.