## For Research Use Only

# TDP-43 Monoclonal antibody

Catalog Number:66734-1-lg 4 Publications



**Basic Information** 

Catalog Number: 66734-1-lg Size:

1500 µg/ml Source: Mouse Isotype: lgG1

Full Name: TAR DNA binding protein Calculated MW:

> 43 kDa Observed MW: 43 kDa

BC001487

23435

Q13148

GeneID (NCBI):

**UNIPROT ID:** 

GenBank Accession Number:

**Purification Method:** 

1G4F5

Protein A purification CloneNo.:

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000

**Applications** 

**Tested Applications:** FC, IHC, WB, ELISA Cited Applications: IF, IHC, WB Species Specificity:

Human, Rat, Mouse **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: HeLa cells, HEK-293 cells, Jurkat cells, HSC-T6

cells, C6 cells, NIH/3T3 cells

IHC: mouse brain tissue, rat liver tissue, human gliomas tissue

# **Background Information**

Transactivation response (TAR), DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43), was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major  $component \ of \ the \ tau-negative \ and \ ubiquitin-positive \ inclusions \ that \ characterize \ amyotrophic \ lateral \ sclerosis$ (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). 66734-1-Ig is a mouse monoclonal antibody recognizing the intact protein of TDP-43.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Phaneendra Chennampally	34518579	Sci Rep	IF
Rui Zhang	33613100	Int J Biol Sci	WB, IF
Alissa L Nana	30511086	Acta Neuropathol	IF

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 66734-1-1g (TDP-43 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66734-1-Ig (TDP-43 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66734-1-Ig (TDP-43 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 SH-SY5Y cells were intracellularly stained with 0.20 ug/test Anti-Human TDP-43 (66734-1-lg, Clone:1G4F5) (red) or 0.20 ug control antibody (blue) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) with dilution 1:1000. Fixed with90% MeOH.