

For Research Use Only

PPARA Monoclonal antibody

Catalog Number: 66826-1-Ig

Featured Product

49 Publications



Basic Information

Catalog Number:

66826-1-Ig

Size:

1000 μ g/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG7896

GenBank Accession Number:

BC000052

GeneID (NCBI):

5465

UNIPROT ID:

Q07869

Full Name:

peroxisome proliferator-activated receptor alpha

Calculated MW:

52 kDa

Observed MW:

53 kDa

Purification Method:

Protein A purification

CloneNo.:

1G1E10

Recommended Dilutions:

WB 1:1000-1:6000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

Human, rat

Cited Species:

human, rat, mouse, pig

Positive Controls:

WB : HSC-T6 cells, ROS1728 cells

Background Information

Peroxisome proliferator-activated receptor alpha (PPARA) is a ligand-activated transcription factor that belongs to the PPAR nuclear receptor superfamily. PPARA is essential in the modulation of lipid transport and metabolism, mainly through activating mitochondrial and peroxisomal fatty acid β -oxidation pathways. In addition, PPARA seems to decrease inflammation mainly through direct interaction with NF- κ B, causing inhibition of its signaling pathway or reducing the activated levels of NF- κ B and subsequent inflammation. Furthermore, PPARA was implicated in the attenuation of oxidative stress in alcoholic liver disease when treated with polyene phosphatidylcholine through downregulation of ROS-generating enzymes such as ethanol-inducible cytochrome P450 2E1 (CYP2E1), acyl-CoA oxidase, and NADPH oxidase. PPARA exists two isoforms and molecular weight of PPARA isoforms are 52 kDa and 22 kDa. The ability of a retinoid X receptor (RXR) to heterodimerize with many nuclear receptors, including LXR, PPAR, NGF1B and RAR, underscores its pivotal role within the nuclear receptor superfamily. Among these heterodimers, PPAR:RXR is considered an important signalling mediator of both PPAR ligands, such as fatty acids, and 9-cis retinoic acid (9-cis RA), an RXR ligand. (PMID: 15103326). PPARA can form Heterodimer with RXRA and molecular weight of Heterodimer is about 110 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Jia Xu	36210393	Fish Physiol Biochem	WB
Zhonghao Li	36498935	Int J Mol Sci	WB
Xin Yin	35534547	Cell Death Differ	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 66826-1-Ig (PPARA antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.